



ZDC - EIC

Hsin-Yi Chou

Institute of Physics, Academia Sinica





Dec 06, 2024

AMS Electromagnetic Calorimeter (ECAL)



Direction Projection of Layers (10 in Z-Y, 8 in Z-X)

Y	Y	X	X	Y	Y	X	X	Y	Y	X	X	Y	Y	X	X	Y	Y
---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---



The longitudinal shower profile is described by the Gamma distribution:

$$\frac{dE}{dt}(t) = E_0 \frac{(\beta t)^{\beta T_0} \beta e^{-\beta t}}{\Gamma(\beta T_0 + 1)}, \quad (1)$$

using parameters described above. The scale parameter β is found to be constant over the wide energy range, therefore it is fixed in the present analysis. Individual shower parameters E_0 and T_0 are obtained from a fit to observed energy depositions in the ECAL cells of that shower. Fig. 7 shows the high quality of the description of electron showers over a wide energy range by the model based on Eq. (1).

At a given shower depth, the transverse shower shape as a function of the distance from the shower axis r is described by the sum of a narrow core and a wide tail:

$$\frac{dE}{dr}(t, r) \propto Q_C \frac{2rR_C^2}{(r^2 + R_C^2)^2} + (1 - Q_C) \frac{2rR_T^2}{(r^2 + R_T^2)^2}, \quad (2)$$

