Science with the Simons Observatory Large Aperture Telescope



Colin Hill (Columbia)

on behalf of the Simons Observatory Collaboration

New Frontiers in Observational Cosmology with Gravitational Waves and Cosmic Microwave Background NCU, Taipei 7 November 2025

arXiv: 1808.07445 2503.00636


















The Simons Observatory is funded by generous grants from the Simons Foundation and the Heising-Simons Foundation



as well as major contributions from the Founding Institutions, the US NSF, UKRI, and Japan














Colin Hill Simons Observatory Science Goals and Probes Columbia



10

Additional science includes (but is not limited to):

- · helium fraction, cosmic birefringence, primordial magnetic fields
- high-redshift clusters
- dark matter annihilation and interactions
- isocurvature
- calibration of multiplicative shear bias (e.g., for LSST)

100

- new sample of dusty star-forming galaxies
- transient sources
- cosmic infrared background

1000


THE SIMONS OBSERVATORY: SCIENCE GOALS

AND FORECASTS

1808.07445 2503.00636




Simons Observatory Site





Colin Hill Simons Observatory Instruments & Technology Columbia

large aperture telescope **15m** ~30,000 detectors 6 m crossed Dragone fed by up to 13, 38 cm optics tubes. baseline=7 tubes for SO, with baseline pixels: One tube: 30/40 GHz Four tubes: 90/150 GHz Two tubes: 220/270 GHz




+ 3 additional SATs coming online by 2027

+ 6 additional OTs installed in 2026

~60,000 detectors in total!



Simons Observatory LAT Timeline





Anticipated Noise Performance

Colin Hill Columbia

SO LAT: 2025-2034

assuming 6 additional optics tubes are added in 2028

Frequency	FWHM	Baseline Depth	Goal Depth	Frequency	Detector	Optics
[GHz]	[arcmin]	$[\mu K \cdot arcmin]$	$[\mu K \cdot arcmin]$	Bands	Count	Tubes
$27\ (22-30)$	7.4	61	44	LF	354	1
39 (30-47)	5.1	30	23	Γ	354	1
$93\ (77-104)$	2.2	5.3	3.8	MF	20,640	Q
145(128-169)	1.4	6.6	4.1	IVIT	20,640	O
$225 \ (198 - 256)$	1.0	15	10	UHF	10,320	1
$280\ (256-313)$	0.9	35	25		10,320	4

Combined sensitivity (goal): 2.6 µK*arcmin

Significant improvement over previous surveys:

Planck: ~25 µK*arcmin (~5 arcmin resolution)


ACT DR6: ~10 µK*arcmin



Anticipated Noise Performance

Significant gains from NSF-funded additional optics tubes (~40% increase in number of signal-dominated E-modes)








Anticipated Sky Coverage

Colin Hill Columbia



total $f_{sky} \sim 60\%$ (cosmological $f_{sky} \sim 40\%$)

maximal overlap w/ Rubin LSST, large overlap w/ DESI

SO LAT Science



- Extragalactic astrophysics: galaxy formation, ICM physics
- Particle physics: neutrino masses, new light species, BSM
- Primordial perturbations: non-Gaussianity, P(k)
- Sources: variable blazars, dusty galaxies
- Transients: uncharted discovery space
- Milky Way ISM: dust composition, magnetic fields
- Solar System: Planet 9, asteroids
 - + much more


SO LAT Science

- Extragalactic astrophysics: galaxy formation, ICM physics
- Particle physics: neutrino masses, new light species, BSM
- Primordial perturbations: non-Gaussianity, P(k)
- Sources: variable blazars, dusty galaxies
- Transients: uncharted discovery space
- Milky Way ISM: dust composition, magnetic fields
- Solar System: Planet 9, asteroids
 - + much more

Colin Hill
Columbia

Cosmic Microwave Backlight


Secondary Anisotropies



Colin Hill Columbia


Cosmic Microwave Backlight

Secondary Anisotropies



- Deflection: gravitational lensing
- Evolving potentials: integrated Sachs-Wolfe, Rees-Sciama effects
- Scattering: thermal / kinematic Sunyaev-Zel'dovich effects, patchy screening






Kinematic Sunyaev-Zel'dovich Effect:

Doppler boosting of CMB photons
Compton-scattering off free electrons
with non-zero line-of-sight velocity

Direct probe of gas momentum

→ gas abundance





Sudeep Das/ACT

Kinematic Sunyaev-Zel'dovich Effect:

Doppler boosting of CMB photons
Compton-scattering off free electrons
with non-zero line-of-sight velocity


Direct probe of gas momentum


→ gas abundance

Thermal SZ Effect:

Change in temperature of CMB photons due to inverse Compton scattering off hot electrons, most of which are in the intracluster medium (ICM) of galaxy clusters


Direct probe of gas pressure







Velocity-weighted kSZ stacking



- Direct imaging of gas density (kSZ) and pressure (tSZ) in the CGM
- Current stacked kSZ measurements (ACT+BOSS/DESI): S/N~10
- SO stacked kSZ measurements (SO+DESI or Rubin): S/N~100s

Direct constraints on galaxy formation physics from stacked kSZ+tSZ


 η = efficiency of feedback energy injection; P_{nt} = non-thermal pressure support in the CGM/ICM

$$\sigma(\eta) \qquad \sigma(P_{nt})$$
 Current (ACT + BOSS) 25% unconstrained Nominal SO + DESI 3% 5%
$$Full \ SO + Rubin + DESI \ 0.5\%$$
 1%

Direct constraints on sub-grid AGN and supernova feedback parameters in galaxy formation simulations:


Current (ACT + BOSS): unconstrained Nominal SO + DESI: ~7%-30% Full SO + Rubin + DESI: ~2%-10%

Amodeo+21 [ACT]
CAMELS (Moser+22)





First statistical census of galaxy clusters at z > 2




- High-sensitivity, high-resolution tSZ surveys can detect galaxy clusters out to the highest redshifts at which they exist
- Nominal SO: ~40 clusters at z>2 (24,000 total)
- Full SO: ~200 clusters at z>2 (33,000 total)

Hilton+21 [ACT]




First statistical census of galaxy clusters at z > 2



Hilton+21 [ACT]

- High-sensitivity, high-resolution tSZ surveys can detect galaxy clusters out to the highest redshifts at which they exist
- Nominal SO: ~40 clusters at z>2 (24,000 total)
- Full SO: ~200 clusters at z>2 (33,000 total)

Spectral coverage for relativistic tSZ






- Extragalactic astrophysics: galaxy formation, ICM physics
- Particle physics: neutrino masses, new light species, BSM
- Primordial perturbations: non-Gaussianity, P(k)
- Sources: variable blazars, dusty galaxies
- Transients: uncharted discovery space
- Milky Way ISM: dust composition, magnetic fields
- Solar System: Planet 9, asteroids
 - + much more




Primary CMB





CMB: Relativistic Particle Detector



- The CMB power spectra are sensitive to any light particles (mass < eV) that were ever in thermal contact with the primordial plasma ("dark radiation", e.g., neutrinos)
- N_{eff} is a simple parametrization that captures a wide range of beyond-standard-model theories, with canonical value N_{eff} = 3.044 for the SM neutrinos:

$$N_{\mathrm{eff}} = \frac{8}{7} \left(\frac{11}{4}\right)^{4/3} \frac{\rho_{\nu}}{\rho_{\gamma}}$$


Current Data [ACT+Planck] SO Nominal			Full SO	
$\sigma(N_{_{ m eff}})$	0.13	0.06	0.045	

SO: sensitive to any light spin-3/2 particle that was in thermal contact at any time back to reheating




CMB: Potential Hints of New Physics in H₀

- Disagreement has reached 5σ between CMB and SH0ES: if not a systematic (robust confirmation still needed), this requires new physics beyond ΛCDM
- Not a CMB systematic: Planck/ACT/SPT agree within ΛCDM
- Not specific to CMB: other cosmological probes (BAO + BBN + others) agree
- How can we increase the value of H₀ inferred from CMB and large-scale structure data?
- → Introduce new physics just prior to recombination that modifies the physical size of the sound horizon
- → This yields novel signals in the CMB power spectra



CMB: Potential Hints of New Physics in H₀


- Disagreement has reached 5σ between CMB and SH0ES: if not a systematic (robust confirmation still needed), this requires new physics beyond ΛCDM
- Not a CMB systematic: Planck/ACT/SPT agree within ΛCDM
- Not specific to CMB: other cosmological probes (BAO + BBN + others) agree
- How can we increase the value of H₀ inferred from CMB and large-scale structure data?
- → Introduce new physics just prior to recombination that modifies the physical size of the sound horizon
- → This yields novel signals in the CMB power spectra



Many well-motivated models now exist with the potential to explain the H₀ problem via new physics that can only be seen unambiguously via high-S/N measurements of the CMB polarization power spectra

- Early dark energy (Poulin+19, Lin+19, Agrawal+19, Hill+21) / early dark sector (McDonough, Lin, Hill+21)
- Strongly interacting neutrinos (Kreisch+19)
- "Stepped" strongly interacting dark radiation (Aloni+21)
- Mirror world dark sector (Cyr-Racine+21)
- Primordial magnetic fields (Jedamzik & Pogosian 19)
- Varying electron mass + curvature (Sekiguchi & Takahashi 20)

CMB: Potential Hints of New Physics in H₀



Example: early dark energy (EDE) scenario

Idea: brief period of cosmic acceleration just prior to recombination

Example EDE and Λ CDM models

EDE ($f_{EDE} \sim 0.2$) - Planck best-fit Λ CDM EDE ($f_{EDE} \sim 0.1$) - Planck best-fit Λ CDM

Key differences are seen in pol. power spectra – but signal is tiny

We need as many high-S/N modes as possible to maximize discovery potential – low noise and wide area:

If EDE were present: 50σ in full SO data

Hill et al. [ACT] (2021)



Neutrino Mass... Need τ!

Astro2020: "[constraining neutrino masses] is typically done by comparing the amplitude of clustering at z = 1100 from CMB observations to that in the late-time evolved universe."

origin of the τ degeneracy


Target signal: 0.06 eV (minimal total mass allowed by neutrino oscillation data)



Neutrino Mass... Need τ!

Epoch of Reionization

Reionization optical depth τ is by far the least well-constrained parameter of ΛCDM: SO can improve upon current constraints by factor of 2.5 using kinematic SZ data



Planck E-mode pol.: $\sigma(\tau) \sim 0.1$

Nominal SO kSZ: $\sigma(\tau) = 0.007$

Full SO kSZ: $\sigma(\tau) = 0.0035$

Alvarez, Ferraro, Hill, Hlozek, Ikape 20; Smith & Ferraro 17,18; Calabrese+14;

Neutrino Mass

Astro2020: "[constraining neutrino masses] is typically done by comparing the amplitude of clustering at z = 1100 from CMB observations to that in the late-time evolved universe."

origin of the τ degeneracy

Target signal: 0.06 eV (minimal total mass allowed by neutrino oscillation data)

SO will use multiple probes to constrain M_v via the late-time amplitude of structure (similar to dark energy)

Focus here on the CMB lensing power spectrum + primary CMB + τ from reionization kSZ 2pt/4pt

	Current Data	SO Nominal [w/ kSZ τ]	Full SO [w/ kSZ τ]
σ(M _v) [eV]	0.05	0.03	0.02 [> 3σ detection]

No external data assumed here (LiteBIRD/CLASS)


(0.03 eV, with Planck EE τ) (0.015 eV, with CV-limited EE τ from external data)



Dark Energy

Constraints derived from tSZ cluster counts (+LSST WL and/or SO CMB lensing), and CMB lensing cross-correlations w/ LSST galaxies

Amplitude of fluctuations as a function of redshift



DE equation of state (via tSZ cluster counts)

$$\sigma(w_0) = 0.06, \quad \Lambda \text{CDM} + w_0 + w_a \ \sigma(w_a) = 0.20, \ \sigma(w_0) = 0.08, \quad \Lambda \text{CDM} + w_0 + w_a + \Sigma m_{\nu} \ \sigma(w_a) = 0.32,$$

Unique ability to search for deviations from Λ at z > 1 (complementary to lower-redshift WL and galaxy surveys)

(N.B. non-red lines are nominal-SO errors)


SO LAT Science

- Extragalactic astrophysics: galaxy formation, ICM physics
- Particle physics: neutrino masses, new light species, BSM
- Primordial perturbations: non-Gaussianity, P(k)
- Sources: variable blazars, dusty galaxies
- Transients: uncharted discovery space
- Milky Way ISM: dust composition, magnetic fields
- Solar System: Planet 9, asteroids
 - + much more



Primordial Scalar Perturbations

Primordial scalar power spectrum



(N.B. these are nominal-SO errors)


$$\sigma(n_s) = 0.002$$






Primordial Scalar Perturbations

Primordial scalar power spectrum



Local-type primordial non-Gaussianity



(N.B. these are nominal-SO errors) =

Shape $(\langle \zeta \zeta \zeta \rangle)$ $\langle TTT \rangle, \langle TTE \rangle,$ $\langle TEE \rangle, \langle EEE \rangle$	Current (Planck)	SO Baseline	SO Goal
local	5	4	3
equilateral	43	27	24
$\overline{\text{orthogonal}}$	21	14	13
~ //			


+ tensor NGs






Primordial Scalar Perturbations

Primordial scalar power spectrum



(N.B. these are nominal-SO errors)

Local-type primordial non-Gaussianity



can reach $\sigma(f_{NL})$ < 1 with sufficient control of large-scale systematics in SO CMB lensing x LSST galaxies


SO LAT Science



- Extragalactic astrophysics: galaxy formation, ICM physics
- Particle physics: neutrino masses, new light species, BSM
- Primordial perturbations: non-Gaussianity, P(k)
- Sources: variable blazars, dusty galaxies
- Transients: uncharted discovery space
- Milky Way ISM: dust composition, magnetic fields
- Solar System: Planet 9, asteroids
 - + much more



Source Sensitivity



SO will detect ~96,000 AGN and ~30-40,000 dusty galaxies ~daily cadence monitoring for 7,500 of these AGN




may detect ~100 SMBH binaries via sinusoidal light curves

Variable AGN



SO will continuously monitor and publicly release light curve data for ~40,000 AGN (after completion of NSF-funded pipeline in ~2028)

Enables multi-wavelength and multi-messenger (e.g., neutrino) probes of AGN physics





SO LAT Science

- Extragalactic astrophysics: galaxy formation, ICM physics
- Particle physics: neutrino masses, new light species, BSM
- Primordial perturbations: non-Gaussianity, P(k)
- Sources: variable blazars, dusty galaxies
- Transients: uncharted discovery space
- Milky Way ISM: dust composition, magnetic fields
- Solar System: Planet 9, asteroids
 - + much more










Known classes of extragalactic synchrotron transients (100 GHz)





minimum
volumetric event
rate leading to one
transient detection


Duration


Eftekhari+ 2021

Known classes of









We estimate characteristic luminosities and intrinsic rates

Duration


Eftekhari+ 2021



Transients

Colin Hill Columbia

SO will be sensitive to known classes of millimeter-wave transients and to new discoveries




Detection Distance [Mpc]

Transients

SO will probe the physics of jet launching and collimation by detecting GRB reverse shocks.

- SO will provide the first unbiased survey of millimeter-wave GRB reverse shock emission. SO 's large sample of reverse shock emission will enable detailed study of jet launching and collimation.
- SO polarization measurements will constrain jet magnetization.





Transients

Estimated detection rates (some are highly uncertain)


Class	Volumetric Rate	$L_ u$	Distance	Detection Rate
	$(\mathrm{yr}^{-1}\mathrm{Mpc}^{-3})$	$({\rm erg s^{-1} Hz^{-1}})$	(Mpc)	
Extragalactic fast (1–10 d duration)	_	_	_	< 10
Long-duration GRB, on-axis	4×10^{-10}	10^{32}	1300	2
Long-duration GRB, off-axis	6×10^{-9}	3×10^{30}	360	0.6
TDE, on-axis	3×10^{-11}	10^{31}	670	0.02
Low-luminosity GRB	2×10^{-7}	$10^{29} – 10^{30}$	70 – 210	0.1 - 4
AT2018cow-like FBOT	10^{-7}	10^{30}	210	2
Interacting SN	10^{-8}	$10^{27} - 10^{30}$	< 210	$\lesssim 0.2$
Stellar flares	_	_	_	180


Table 4. Estimated detection rates by SO for different classes of extragalactic transients at 145 GHz, and for stellar flares at 90 GHz. We define off-axis long-duration GRBs to have $\theta_{\rm obs} = 0.4$.


(GRB reverse shocks not listed here, but will be numerous)

SO LAT Science

- Extragalactic astrophysics: galaxy formation, ICM physics
- Particle physics: neutrino masses, new light species, BSM
- Primordial perturbations: non-Gaussianity, P(k)
- Sources: variable blazars, dusty galaxies
- Transients: uncharted discovery space
- Milky Way ISM: dust composition, magnetic fields
- Solar System: Planet 9, asteroids
 - + much more








Polarized Galactic Emission


SO will make deep maps of polarized Galactic emission ACT total intensity + 90 GHz polarization





Polarized Galactic Emission Magnetic fields in molecular clouds

- The role of magnetic fields in the formation and evolution of molecular clouds is poorly understood.
- SO will measure magnetic fields at scales intermediate between *Planck* and ALMA
- 0.9' angular resolution at 280 GHz corresponds to 0.03 pc at a distance of 100 pc
- SO will measure the magnetic field structure of >860 molecular clouds with 1 pc resolution and at least 50 independent, high-SNR polarization measurements per cloud






Polarized Galactic Emission

The composition of interstellar dust

- Does interstellar dust consist of distinct silicate and carbonaceous grains?
- We will test and constrain the classical dust models that predict a divergence of the total intensity and polarization spectra in the frequency range probed by SO.
- We will further constrain dust grain shape and porosity via comparison to starlight polarization measurements

Constraint on dust spectral index: $\sigma(\beta_{dust}) = 0.005$




Hensley & Draine 2021






Probing solar system formation via exo-Oort clouds

- The Oort Cloud is a roughly isotropic shell of ice and dust that is thought to surround our Sun at R~10³
 10⁵ AU
- We infer the presence of the Oort Cloud dynamically, but it has never been directly observed.
- SO will search for thermal emission from exo-Oort clouds around nearby Gaia stars





SO LAT Science




- Extragalactic astrophysics: galaxy formation, ICM physics
- Particle physics: neutrino masses, new light species, BSM
- Primordial perturbations: non-Gaussianity, P(k)
- Sources: variable blazars, dusty galaxies
- Transients: uncharted discovery space
- Milky Way ISM: dust composition, magnetic fields
- Solar System: Planet 9, asteroids
 - + much more



Solar System


SO can search for thermal emission from distant Solar System objects (e.g., "Planet 9")





Solar System

SO can search for thermal emission from distant Solar System objects (e.g., "Planet 9")




- SO can detect a 5 Earth-mass planet out to ~900 AU in the deepest regions of the survey (~500 AU in shallowest)
- + 10⁴ asteroids and many of the largest TNOs (e.g., Pluto)

SO LAT: Status













SO LAT Science: Summary














https://arxiv.org/abs/1808.07445 https://arxiv.org/abs/2503.00636



Thanks!

https://simonsobservatory.org/



SO LAT Science: Summary

SO Covers All of A2020/P5 Science

Search for B-modes from the infant universe & inflation Study the dark universe signatures

Matter mapping

	v	v	
Numbers show 1-σ unless	Current	SO	Using Rubin,
catalog no. or distances		2025-2034	DESI, or Euclid
Primordial perturbations			
$r (A_L = 0.3)$	0.01	0.0012	✓
n_s	0.003	0.002	_
$e^{-2 au}\mathcal{P}(k=0.2/\mathrm{Mpc})$	1%	0.4%	_
$f_{ m NL}^{ m local}$	5	1	✓
Relativistic species			
$N_{ m eff.}$	0.13	0.045	-
Neutrino mass			
$m_{\nu} \; (\text{eV}, \sigma(\tau) = 0.01)$	0.06	0.03	✓
$m_{\nu} \; (\text{eV}, \sigma(\tau) = 0.002)$		0.015	✓
Accelerated expansion			
$\sigma_8(z=1-2)$	7%	1%	✓
Galaxy evolution			
$\eta_{ m feedback}$	50-100%	2%	✓
$p_{ m nt}$	50-100%	4%	✓
Reionization			
Δz	~1.4	0.3	-
au	0.007	0.0035	-
Cluster catalog	10000	33,000	✓
AGN catalog	20000	96,000	-
Galactic science			
Molecular cloud B-fields	10s	> 860	-
$\sigma(eta_{dust})$	0.02	0.005	-
Solar System Science			
Distance limit for 5 M_{\oplus} Planet 9	500 AU	900 AU	✓
Asteroid detections		$\sim 10,000$	
Transient detection			
distance			
Long GRBs, on-axis		1300 Mpc	-
Low-luminosity GRBs		70–210 Mpc	-
TDEs, on-axis		670 Mpc	-

https://arxiv.org/abs/1808.07445 https://arxiv.org/abs/2503.00636 & SAT white paper, in prep

Time-variable mm sky