

Moments of pion distribution amplitude using OPE on lattice

Santanu Mondal*

Collaborators: William Detmold**, Issaku Kanamori\$, C.-J.
David Lin* and Yong Zhao**

Academia Sinica

28th June, 2018

*Institute of Physics, National Chiao-Tung University, Taiwan

**Centre for Theoretical Physics, Massachusetts Institute of Technology

\$ Hiroshima University

Outline

- Pion distribution amplitude and its moments
- Moments using lattice OPE with a valance heavy quark ¹
- Lattice correlators
- Exploratory numerical results
- Summary

¹proposed by Detmold and Lin (Phys.Rev. D73 (2006)). A set of small, light-blue navigation icons typically used in Beamer presentations for navigating between slides and sections.

Pion LC wave function/distribution amplitude:

$$\langle \pi^+(p) | \bar{d}(\frac{z}{2}) \gamma_5 \gamma_\mu u(-\frac{z}{2}) | 0 \rangle = -ip_\mu f_\pi \int_0^1 d\xi e^{i(\bar{\xi} p \frac{z}{2} - \xi p \frac{z}{2})} \phi_\pi(\xi)$$
$$\bar{\xi} = 1 - \xi$$

fraction ξ of pion momentum is carried by u quark.

(Mellin) Moments:

$$a_n = \int_0^1 d\xi \xi^n \phi_\pi(\xi).$$

OPE:

$$\langle \pi^+(p) | O^{\mu_1 \dots \mu_n} | 0 \rangle = f_\pi a_{n-1} [p^{\mu_1} \dots p^{\mu_n} - \text{Traces}]$$
$$O^{\mu_1 \dots \mu_n} = \bar{\psi} \gamma^{\{\mu_1} \gamma^5 (iD^{\mu_2}) \dots (iD^{\mu_n\}}) \psi - \text{Traces}$$

In the isospin limit $m_u = m_d$:

$$\phi_\pi(\xi) = \phi_\pi(\bar{\xi})$$

⇒ Odd moments vanish → lowest non-trivial moment is a_2 .

Lattice calculation:

- second moment is calculated quite precisely ([Braun et. al., 2015](#)).
- going beyond is challenging because:
 - operator mixing (reduced symmetry: $O(4) \rightarrow H(4)$).

A different approach is needed.

Euclidean OPE with a valance heavy quark

We need to calculate the matrix elements:

$$\int d^4x e^{iqx} \langle \pi^+(p) | T[A_{\Psi,\psi}^\mu(x) A_{\Psi,\psi}^\nu(0)] | 0 \rangle$$

Heavy-light currents:

$$A_{\Psi,\psi}^\mu = \bar{\Psi} \gamma^\mu \gamma^5 \psi + \bar{\psi} \gamma^\mu \gamma^5 \Psi$$

ψ : light quarks, Ψ : fictitious, relativistic, valance quark which is heavy.


- Simplify the lattice calculation:
 - no disconnected contribution.
 - easy to invert the quark matrix.
- No effect of the heavy quark on DA.

Scale hierarchy required:

$$\Lambda_{QCD} \ll m_\Psi \ll \frac{1}{a}$$

⇒ Fine lattices are required.

Also, need to take the continuum limit.



VV type operator:

$$\int d^4x e^{iqx} T[A_{\Psi,\psi}^\mu(x)A_{\Psi,\psi}^\nu(0)] = \sum_k C_k(q) \mathcal{O}_k(0) \text{ for large } q$$

Euclidean OPE:

$$\int d^4x e^{iqx} T[A_{\Psi,\psi}^\mu(x)A_{\Psi,\psi}^\nu(0)] = \bar{\psi} \gamma^\mu \frac{-i(iD + q) + m_\psi}{(iD + q)^2 + m_\psi^2} \gamma^\nu \psi + \bar{\psi} \gamma^\nu \frac{-i(iD - q) + m_\psi}{(iD - q)^2 + m_\psi^2} \gamma^\mu \psi$$

Taylor expansion:

$$\frac{-i(iD + q) + m_\psi}{(iD - q)^2 + m_\psi^2} = -\frac{-i(iD + q) + m_\psi}{Q^2 + D^2 - m_\psi^2} \sum_{n=0}^{\infty} \left(\frac{-2iq.D}{Q^2 + D^2 - m_\psi^2} \right)^n, \quad Q^2 = -q^2$$

Higher twist terms \rightarrow expansion parameter: $\left(\frac{-2iq.D+D^2}{Q^2-m_\Psi^2}\right) \rightarrow$ Extra powers of $p^2 \rightarrow$ small.

Antisymmetric in μ and ν :

$$\int d^4x e^{iqx} T[A_{\Psi,\psi}^\mu(x) A_{\Psi,\psi}^\nu(0)] = \frac{i}{2} \sum_{n=0, \text{even}}^{\infty} \frac{1}{(\tilde{Q}^2)^{n+1}} \bar{\psi} \gamma_\lambda \gamma_5 (i \not{D}_{\mu_1}) \dots (i \not{D}_{\mu_n}) \psi \epsilon_{\mu\nu\rho\lambda}$$

$$\tilde{Q}^2 = Q^2 - m_\Psi^2$$

$$U_A^{[\mu\nu]}(p,q) = \int d^4x e^{iqx} \langle \pi^+(p) | T[A_{\Psi,\psi}^\mu(x) A_{\Psi,\psi}^\nu(0)] | 0 \rangle$$

$$= f_\pi \epsilon_{\mu\nu\rho\lambda} q^\rho p^\lambda \sum_{n=0, \text{even}}^{\infty} a_n f_n(q^2, p.q, m_\Psi)$$


$$f(n) = \frac{i}{2} \frac{\xi^{n+1}}{n+1} \left[\frac{2\eta C_n^2(\eta)}{p.q} \right], \xi = \frac{\sqrt{p^2 q^2}}{\tilde{Q}^2}, \eta = \frac{p.q}{\sqrt{p^2 q^2}}$$

↑

$$f(n) \sim \frac{1}{\xi^n} \sim \left(\frac{1}{\tilde{Q}^2}\right)^{(n+1)}$$

- Analytic continuation is in the Wilson coefficients.
- For simplicity, Wilson coefficients are set to one.
- Identical result for the VV type correlator.

Lattice correlators: the three point-function



$$C_3^{\mu\nu}(\tau_m, \tau_e; \vec{p}_m, \vec{p}_e) = \int d^3x_m \int d^3x_e e^{i\vec{p}_e \cdot \vec{x}_e} e^{-i\vec{p}_m \cdot \vec{x}_m} \langle 0 | T[A_e^\mu(\vec{x}_e, \tau_e) A_m^\nu(\vec{x}_m, \tau_m) \mathcal{O}_\pi^\dagger(\vec{0}, 0)] | 0 \rangle$$

Two different choices:

- shift the operator $T[A_e^\mu(\vec{x}_e, \tau_e)A_m^\nu(\vec{x}_m, \tau_m)]$ by (\vec{x}_m, τ_m) .
- shift the operator $T[A_e^\mu(\vec{x}_e, \tau_e)A_m^\nu(\vec{x}_m, \tau_m)]$ by (\vec{x}_e, τ_e) .

$$\begin{aligned} C_3^{\mu\nu}(\tau_m, \tau_e; \vec{p}_m, \vec{p}_e) &= \frac{1}{2V_3 E_\pi} \delta_{\vec{n}, \vec{n}_e - \vec{n}_m} \times e^{-E_\pi \tau_m} \langle \pi(\vec{p}) | \mathcal{O}_\pi^\dagger(\vec{0}, 0) | 0 \rangle \\ &\quad \times \int d^3x e^{i\vec{p}_e \cdot \vec{x}} \langle 0 | T[A_e^\mu(\vec{x}, \tau) A_m^\nu(\vec{0}, 0)] | \pi(\vec{p}) \rangle \\ &\quad \vec{x} = \vec{x}_e - \vec{x}_m, \tau = \tau_e - \tau_m : \end{aligned}$$

The two point function:

$$\begin{aligned} C_\pi(\tau_\pi; \vec{p}_\pi) &= \int d^3x e^{i\vec{p}_\pi \cdot \vec{x}} \left\langle 0 \left| \mathcal{O}_\pi(\vec{x}, \tau) \mathcal{O}_\pi^\dagger(\vec{0}, 0) \right| 0 \right\rangle \\ &\xrightarrow{\tau_\pi \rightarrow \infty} \frac{\left| \left\langle \pi(\vec{p}_\pi) \left| \mathcal{O}_\pi^\dagger(\vec{0}, 0) \right| 0 \right\rangle \right|^2}{2V_3 E_\pi} \times e^{-E_\pi \tau_\pi}. \end{aligned}$$

We can form the ratio

$$\begin{aligned} R_{3,m}^{\mu\nu}(\tau; \vec{q}, \vec{p}) &= \frac{C_3^{\mu\nu}(\tau_m, \tau_e; \vec{p}_e, \vec{p}_m)}{C_\pi(\tau_m; \vec{p})} \times \left\langle \pi(\vec{p}_\pi) \middle| \mathcal{O}_\pi^\dagger(\vec{0}, 0) \middle| 0 \right\rangle \\ &= \int d^3x e^{i\vec{q}\cdot\vec{x}} \left\langle 0 \middle| T[A_e^\mu(\vec{x}, \tau) A_m^\nu(\vec{0}, 0)] \middle| \pi(\vec{p}_\pi) \right\rangle_{\tau=\tau_e-\tau_m; \vec{q}=\vec{p}_e; \vec{p}=\vec{p}_e-\vec{p}_m} \end{aligned}$$

Perform the temporal Fourier transform:

$$\int d\tau e^{iq_4\tau} R_{3,m/e}^{\mu\nu}(\tau; \vec{q}, \vec{p}) = U_A^{\mu\nu}(q, p)$$

exploratory numerical results

Quenched calculation:

- Hugely reduces the computational cost.
- Neglecting the effect of the fermion loops.

Nonperturbative Clover action for valance quarks. \rightarrow Wilson quarks with less cut-off artifacts ($\mathcal{O}(a^2)$).


Moderate light quark mass with $m_\pi = 450$ MeV.

Fine lattice: $a^{-1} = 4$ GeV (lattices upto $a^{-1} = 8$ GeV are available).

Moderate physical box size: $L = 2.4$ fm, $T = 4.8$ fm.


Translation invariance:

$$R_3^{[\mu\nu]}(\tau, q, p) + e^{-E_\pi(p)\tau} R_3^{[\mu\nu]}(\tau, p-q, p) = 0$$



$$m_\Psi \sim 2 \text{ GeV} \quad U_A^{[12]} = f_\pi \frac{q^3 p^4 - p^3 q^4}{2} \sum_{n=0, \text{ even}}^\infty a_n f_n(q^2, p \cdot q, m_\Psi)$$


$q=(0,0,1)$, # of configs=226



All have same $p^3 q^4 - q^3 p^4$, $p \cdot q$ and q^2 .


$$m_\Psi \sim 1.3 \text{ GeV}, U_A^{[12]}(q, p) \sim (q^3 p^4 - p^3 q^4)$$

$p=(0,0,0)$, $q=(0,0,1)$, # of configs = 226



$$m_\Psi \sim 1.3 \text{ GeV}, U_A^{[12]}(q, p) \sim (q^3 p^4 - p^3 q^4)$$

$p=(0,0,1)$, $q=(0,0,1)$, # of configs = 226



Summary

- I have described a new method to calculate the moments of pion distribution amplitude by studying the Euclidean OPE on lattice. The method is free from operator mixing problem of the traditional approach.
- A valance heavy quark is used to make lattice calculation simpler and to give more flexibility.
- I have shown our preliminary numerical data and demonstrate our strategy to analyze those. The data is qualitatively consistent with OPE.
- The method described here can also be used to study parton distribution function of the pion and nucleon.

Thanks for your attention!

Lattice Gauge Theory

In QCD at low energy coupling is too strong to investigate with perturbative techniques.

First principle calculation based on path integral formalism.



Path integral in Euclidean space:

$$\langle \mathcal{O}[\bar{\psi}, \psi, A_\mu] \rangle = \frac{1}{Z} \int \mathcal{D}A_\mu \mathcal{D}\bar{\psi} \mathcal{D}\psi \mathcal{O}[\bar{\psi}, \psi, A_\mu] e^{-S[\psi, \bar{\psi}, A_\mu]}$$

Functional integral over continuous space-time of infinite extent
⇒ infinite number of dof's.

Discretize 4-dim. space-time for finite degree of freedom \Rightarrow 4-dim. lattice:

Ultraviolet regulator with cutoff scale $\sim \frac{1}{a}$, a is the lattice spacing.



$$\begin{array}{ccc} \psi(x) & \longrightarrow & \psi(n) \\ A_\mu(x) & \longrightarrow & U_\mu(n), \text{ SU}(3) \text{ group valued link variable} \end{array}$$

Path integral on 4-dim. lattice:

$$\langle \mathcal{O}[\bar{\psi}, \psi, U_\mu] \rangle = \frac{1}{Z} \int \prod_{n,\mu} dU_\mu(n) \bar{\psi}(n) \psi(n) \mathcal{O}[\bar{\psi}, \psi, U_\mu] e^{-\int d^4x \mathcal{L}_{lattice}[\psi, \bar{\psi}, U]}$$

Quark fields are Grassmann (anticommuting) numbers \Rightarrow Integrated analytically

$$\langle \bar{\mathcal{O}}[U_\mu] \rangle = \frac{1}{Z} \int \prod_{n,\mu} dU_\mu(n) \bar{\mathcal{O}}[U_\mu] e^{-S_{lattice}^{eff}[U]}$$

Average over the configurations after importance sampling gives expectation value:

$$\begin{aligned} \langle \bar{\mathcal{O}}[U_\mu] \rangle &= \frac{1}{N} \sum_{i=1}^N \bar{\mathcal{O}}[U_\mu^i] + \mathcal{O}\left(\frac{1}{\sqrt{N}}\right) \\ &\Downarrow \\ &\text{statistical error} \end{aligned}$$

Hadron mass using lattice gauge theory

In continuum notation:

$$\langle \mathcal{O}(x, t) \mathcal{O}^\dagger(0) \rangle = \frac{1}{Z} \int [d\psi][d\bar{\psi}][dU] \mathcal{O}(x, t) \mathcal{O}^\dagger(0) e^{-S[\psi, \bar{\psi}, A_\mu]}$$

Euclidean correlation function:

$$\begin{aligned} C(t) &= \int d^3x \langle \mathcal{O}(x, t) \mathcal{O}^\dagger(0) \rangle \\ &= \int d^3x \langle e^{Ht} e^{i\hat{P}x} \mathcal{O}(0) e^{-Ht} e^{-i\hat{P}x} \mathcal{O}^\dagger(0) \rangle \\ &= \int d^3x \int d^3p \sum_n \langle e^{Ht} e^{i\hat{P}x} \mathcal{O}(0) e^{-Ht} e^{-i\hat{P}x} |E_n, p\rangle \langle E_n, p| \mathcal{O}^\dagger(0) \rangle \\ &= \int d^3x \int d^3p \sum_n \langle \mathcal{O}(0) |E_n, p\rangle \langle E_n, p| \mathcal{O}^\dagger(0) \rangle e^{-E_n t} e^{-ip.x} \\ &= \sum_n |\langle 0 | \mathcal{O}^\dagger(0) |E_n, 0\rangle|^2 e^{-M_n t} \end{aligned}$$

M_n can be extracted by fitting lattice data for $C(t)$.