

# *Activities of the EIC-Japan Group*

NCU Workshop on EIC physics and detectors


NCU, Taoyuan, Taiwan

December 9<sup>th</sup>, 2022

Yuji Goto (RIKEN)

# Electron-Ion Collider (EIC)

- 2020.1.9: U.S. Department of Energy selected Brookhaven National Laboratory to host major new nuclear physics facility, the Electron-Ion Collider
- World's first polarized electron + proton / light-ion / heavy-ion collider



- Center of Mass Energies 20 GeV – 141 GeV
- Maximum Luminosity  $10^{34} \text{ cm}^{-2}\text{s}^{-1}$
- Hadron Beam Polarization 80%
- Electron Beam Polarization 80%
- Ion Species Range p to Uranium
- Number of interaction regions up to two

Polarized beam: e, p, d,  $^3\text{He}$

(Polarized) Ion Source



AGS

- Hadron Storage Ring
- Hadron Injector Complex
- Electron Storage Ring
- Electron Injector Synchrotron
- Electron Cooler
- Possible On-energy Hadron Injector Ring

Electron-Ion Collider 8



# Physics at EIC

- How does the mass of the nucleon arise?
  - The Higgs mechanism accounts for only  $\sim 1\%$  of the mass of the proton.
- How does the spin of the nucleon arise?
  - The spin of the quarks accounts for only one-third of the spin of the proton.
- What are the emergent properties of dense system of gluons?
  - The gluon saturation describes a new state of matter at extreme high density.



# Quark-gluon structure



- Deep inelastic scattering (DIS) of lepton (electron)
  - Large  $Q^2$  ( $Q^2 = -q^2$ ) provides a hard scale to resolve quarks and gluons in the proton
- Parton distribution function (PDF) of quarks and gluons
  - 1D longitudinal motion of partons
  - $x$ : momentum fraction of quarks and gluons
  - Significant improvement of precision of the polarized PDF at EIC







# 3D structure of the nucleon

- Conclusive understanding of the nucleon spin
  - Orbital motion inside the nucleon and orbital angular momenta of quarks and gluons
- TMD (Transverse-Momentum Dependent) distribution function
  - Correlation between the (orbital) motion, spin of partons, and spin of the nucleon






- GPD (Generalized Parton Distribution)
  - Spatial distribution or tomography




# Gluon saturation in $e+A$ collisions

- pQCD and DGLAP & BFKL evolution works with high precision
- Issues with linear DGLAP/BFKL at low- $x$ 
  - Gluon PDF rapid rise violates unitary bound
- New approach: non-linear evolution
  - Gluon emission
    - Divergence at small  $x$
  - Gluon recombination
    - Restriction of divergence
  - At very high energy, recombination compensates gluon emission
- BK/JIMWLK non-linear effects
  - Saturation characterized by  $Q_s(x)$
  - Describe physics at low- $x$  and low-moderate  $Q^2$



# *Gluon saturation in $e+A$ collisions*


- Color Glass Condensate (CGC)
  - Non-linear evolution
  - Saturation of gluon densities characterized by scale  $Q_s(x)$
- Enhancement of  $Q_s$  with  $A$ 
  - Saturation regime reached at significantly lower energy in nuclei



$R \sim A^{1/3}$

$$(Q_s^A)^2 \approx cQ_0^2 \left[ \frac{A}{x} \right]^{1/3}$$

- First observation of a quantum collective gluonic system
  - Precision comparison of experiment and CGC as a theoretical model of the gluon saturation
- Precision understanding of nucleus with the quark-gluon picture necessary as the initial state of the QGP for understanding its production mechanism






# *EIC-Japan activities*

- 2015.4: EIC Letter of Interest from Asian countries
  - 20 participants from Japan: RIKEN, Yamagata, Tokyo Tech, Juntendo, KEK, Kyorin, Kyoto, Niigata, Tohoku, Tokyo Science
  - 7 from China, 3 from India, 4 from Korea
  - To support EIC for NSAC Long Range Plan 2015
- 2019: Science Council of Japan Master Plan 2020 proposal of EIC
  - Collaboration including nuclear-physics community and high-energy community
  - Core institutions: Yamagata and RIKEN
  - Participating institutions: Kobe, Nihon, KEK, etc.
- 2020: Yellow Report
- 2020.5: eRD27 “developing a high resolution ZDC for the EIC”
- 2020.11: Expression of Interest (EOI) from EIC-Japan
- 2021.3-12: Call for detector proposal from the EIC project
  - EIC-Japan group participates in the ECCE detector consortium
- 2022: Science Council of Japan “Medium- and Long-term Research Strategy for Science”
  - EIC project proposal submitted as a part of the High-Energy QCD Frontier Initiative



# *Interest in contributing to ZDC*

- ECCE/EPIC ZDC (Zero-Degree Calorimeter) design
  - Simulation
  - Performance evaluation
- ALICE-FoCal-E technology: Tungsten/Silicon
  - Test beam studies ongoing
- Radiation tolerance test by neutron irradiation
- RIKEN, Tsukuba, Tsukuba Tech, Kobe, Shinshu, Yamagata, JAEA, Nihon, Kyushu, KEK, Nagoya, Tokyo ICRR



**ECCE/EPIC ZDC**


**ALICE FoCal-E  
R&D**

**Neutron irradiation  
at RIKEN RANS**




# *EIC Interaction Region (IP6)*

- Extensive integration of forward and backward detector elements into the accelerator lattice




- EIC far-forward region



# Far-forward physics at EIC


- Spectator tagging in  $e+d/{}^3\text{He}$  collisions

- Neutron structure
  - Neutron spin structure, S & D waves




- $e+A$  collisions at zero degree

- Breakup determination of the excited nucleus
  - Veto with evaporated neutrons and photons from de-excitation
- Geometry tagging in  $e+A$  collisions
  - Event-by-event characterization of collision geometry
  - Study of nuclear medium effects
- Short-range correlation (SRC) and EMC effect
  - Nuclear PDF significantly modified by SRC pairs



Nucleon Momentum Distribution




# Far-forward physics at EIC

- Mass of the proton, pion, kaon
  - Light quarks: its mass emerges from quark-gluon interactions, Higgs mechanism hardly plays a role
  - Strange quark is at the boundary: both emergent-mass and Higgs-mass generation mechanism are important



$$M = E_q + E_g + \chi m_q + T_g$$

X. Ji, PRL 74 1071 (1995)

- Proton
  - Determination of an important term contributing to the proton mass, the so-called “QCD trace anomaly”
  - Through dedicated measurements of exclusive production of  $J/\psi$  and  $\Upsilon$  close to the production threshold
- Pion and kaon
  - Determination of the quark and gluon contribution to mass with the Sullivan process




## Sullivan process Detect scattered electron




# *Interest in contributing to (AC-)LGAD Barrel*

- Construction of (AC-)LGAD (Low-Gain Avalanche Detector) Barrel based on our past experience of PHENIX VTX silicon detector construction and present experience of sPHENIX INTT silicon detector construction
- HPK LGAD development by KEK group
  - To be combined with some readout ASIC
- RIKEN, Hiroshima, Nara Women's, Tokyo CNS, Kyushu, KEK



**sPHENIX INTT construction**



December 9, 2022



**HPK LGAD development**




# Charged particle ID



- Need to separate:
  - Electrons from photons
  - Electrons from charged hadrons
    - Calorimeter
  - Charged pions, kaons and protons from each other
    - TOF and Cherenkov
- AC-LGAD based TOF system
  - Hadron PID in momentum range below the thresholds of the Cherenkov detectors

AC-LGAD TOF (~30 ps)





# *Summary of this talk*

- Physics at EIC
  - Ultra-precise electron microscope, revealing the origin of mass and spin in three dimensions.
  - Discovery of emergent high-density gluon state (gluon condensation)
- EIC-Japan activities
  - Interest in contributing to ZDC
    - Far-forward physics at EIC
  - Interest in contributing to (AC-)LGAD Barrel
    - Charged particle ID
  - EIC-Japan Group is developing steadily

*Backup Slides*

# *Outline of this talk*


- Physics at Electron-Ion Collider (EIC)
  - Origin of nucleon mass and spin
  - 3D structure of the nucleon and nucleus
  - Gluon saturation (Color Glass Condensate)
  - Hadronization
- EIC-Japan activities
  - Interest in contributing to ZDC (Zero-Degree Calorimeter)
  - Interest in contributing to (AC-)LGAD (Low-Gain Avalanche Detector) Barrel
- Collaboration opportunities

# Collaboration opportunities



## • EPIC ZDC

- Soft photon detection
  - Crystal calorimeter (PWO, LYSO, ...)  
prototype
  - Readout device (APD, PMT, ...)
- EM+hadron calorimeter
  - ALICE-FoCal-E technology
    - Pad detector led by Univ. of Tsukuba group and Indian group
    - Pixel detector led by European group
  - Test beam activities ongoing
    - Pad detector at ELPH, Tohoku U.
    - Total system at CERN PS/SPS
  - EM calorimeter optimization
    - Sensor, readout (HGCROC), aggregator
  - Hadron calorimeter design (light collection, readout)

ECCE/EPIC ZDC







ALICE FoCal-E R&D



# EPIC ZDC

- Measurement of the radiation hardness of the ALICE-FoCal-E Pad sensors
  - To determine if the sensor is sufficiently radiation hard to radiation dose/fluence at zero degree of EIC
- Options: p-type or n-type
  - Type inversion from n-type to p-type at  $10^{12}$  neutron/cm<sup>2</sup>





Neutron irradiation at RIKEN RANS



# EPIC ZDC

- At ALICE-FoCal, 1-MeV neutron equivalent fluence  $< 10^{13}$   $n_{eq}/cm^2$ , or Total ionization dose (TID) of 1.5 kGy
- At EPIC ZDC, more than  $2 \times 10^{13}$  neutron/cm<sup>2</sup> in one year at EIC-ZDC
- More than 10 times higher radiation than ALICE-FoCal

Electron-Proton collisions. IP6, p(275)+e(10). Si lifetime in ZDC.



Assume the ep - collision rate is  $1.E+6$  [ep/s]

$$\text{Critical rate } dN_{ep+pg}/dt = 8.E - 1[N/cm^2/ep] * 1.E+6[ep/s] + dN_{pg}/dt =$$

$$= 8.E+5[Hz/cm^2] + .4 E+5[Hz/cm^2]$$

$$\text{ZDC Silicon LifeTime} = 1.E+14 [1/cm^2] / 8.4E+5 [Hz/cm^2] = \sim 0.12 E+9 [s] =$$

$$\sim 4 \text{ years.}$$

# *Collaboration opportunities*

- EPIC ZDC
  - Cooling, support structure
  - Simulation, software development
  - Construction, QA
- EPIC (AC-)LGAD Barrel
  - Yano's talk: Nov. 3 (Thu) afternoon

# *Summary of this talk*

- Physics at EIC
  - Origin of nucleon mass and spin
  - 3D structure of the nucleon and nucleus
  - Gluon saturation (CGC)
  - Hadronization
  - Ultra-precise electron microscope, revealing the origin of mass and spin in three dimensions.
  - Discovery of emergent high-density gluon state (gluon condensation)
- EIC-Japan activities
  - Interest in contributing to ZDC
    - Far-forward physics at EIC
  - Interest in contributing to (AC-)LGAD Barrel
    - Charged particle ID
  - EIC-Japan Group is developing steadily

# *EIC-Japan status*

- Selection is on going for Akiba-san's grant application "Fund for the Promotion of Joint International Research"
- Following this discussion, CNS, U. Tokyo (Gunji) and Hiroshima U. (Shigaki) joined EIC
- Gunji's proposal to the Science Council of Japan (LHC, FAIR, EIC, etc. and Theory) to be a third pillar with J-PARC extension and RIBF upgrade, granted in Japanese Nuclear Physics Committee, and will presented to the Science Council of Japan
  - Koyasu (head of RIKEN Cluster for Pioneering Research) agreed to host this proposal, after communicating with President Gonokami
  - Hatsuda, Saito, Akiba are proposing "RIKEN Platform for international collaborations" which can be an organizational backbone for above proposal
  - Nagamiya agrees to this direction, very positively
  - Gunji and Saito explained the plan to Japan Nuclear Physics Committee in Oct.13, and many members are very positive, but Sakurai was very much against (but the proposal was granted)


# *Science Council of Japan discussion*

- “Medium-and Long-term Research Strategy for Science” for “Future Science Promotion Initiative”
- EIC project proposal to be submitted as a part of the “International High-Energy Quantum Science Frontier: QCD research at overseas facilities”
  - Leading international-collaboration experiments on a long-term basis at overseas facilities
  - Development of common technologies (detector, data processing, and computation technologies)
  - Cooperation between domestic experiment and theory communities
  - Cooperation among experiments according to project trends
  - A major international-collaboration center connecting Japan, Europe, and the U.S.
  - International circulation of cutting-edge knowledge, technology, and people, and human resource development for the next generation



# *International High-Energy Quantum Science Frontier*

- Promote QCD research to be developed at overseas facilities



# *Visit to MEXT*

- 2022.9.13
  - Attendance: Goto, Akiba (RIKEN), Miyachi (Yamagata U.), Shigaki (Hiroshima U.), Yamazaki (Kobe U.)
  - Absence: Gunji (CNS, U. of Tokyo)
- MEXT
  - Office for Particle and Nuclear Research Promotion, Basic and Generic Research Division, Research Promotion Bureau
  - Mr. Takashi Ishikawa (Director)
  - Mr. Atsushi Morikawa (Unit Chief)
  - Ms. Kanako Kumagai (Senior Specialist)
- EIC status
  - EIC project, EIC User Group, EPIC collaboration, EIC-Japan group
  - Resource Review Board
    - MEXT is not yet at the stage where they can participate in the RRB. The situation is uncertain as to how the Science Council of Japan's "Future Science Promotion Initiative" will proceed and how the MEXT will respond to it.
    - Participation in the EIC is an investment in a foreign country and must benefit Japan. We must be able to explain to the Ministry of Finance how the development of this field will lead to the development of academia and the strengthening of national power.
    - If it cannot be explained, it should usually be funded by the Grants-in-Aid for Scientific Research (Kakenhi), the Frontier Budget, or the University Budget.

# *Outline of this talk*

- Physics at EIC
  - Origin of nucleon mass and spin
  - 3D structure of the nucleon and nucleus
  - Gluon saturation
  - Hadronization
- Physics at zero-degree
  - e+A breakup of the excited nucleus & collision geometry
  - e+d/<sup>3</sup>He spectator tagging & short-range correlation
  - pion and kaon structure (and mass)
- EIC status
  - EIC Users Group
  - EIC-Japan
  - EPIC detector collaboration




# *Outline of this talk*

- Physics at Electron-Ion Collider (EIC)
  - Origin of nucleon mass and spin
  - 3D structure of the nucleon and nucleus
  - Gluon saturation (Color Glass Condensate)
  - Hadronization
- EIC status
  - EIC Users Group
  - EIC-Japan
  - EPIC detector collaboration

# Atomic structure

- Scattering experiment of  $\alpha$ -rays
  - $\alpha$ -ray irradiation to gold foil
  - Only small angle scattering if charge is uniformly distributed in atoms (Thomson model)
  - Observation of large angle scattering, discovery of point nuclei, concentration of charge in a narrow region
- Rutherford scattering (1911)

$$\left(\frac{d\sigma}{d\Omega}\right)_{\text{Rutherford}} = \frac{Z^2 \alpha^2}{4E^2 \sin^4 \theta/2}$$



# Structure of nucleus and nucleon

- Electron Beam Scattering Experiment
  - Mott scattering
  - Electron spin 1/2, target recoil

$$\left(\frac{d\sigma}{d\Omega}\right)_{\text{Mott}} = \left(\frac{d\sigma}{d\Omega}\right)_{\text{Rutherford}} \cdot \cos^2 \frac{\theta}{2} \cdot \frac{E'}{E}$$

- Electron-proton elastic scattering
  - Electron beam at SLAC (1950s-60s)
  - Form factor measurement
    - Momentum transfer dependence of angular distribution
  - Rosenbluth formula


$$\frac{d\sigma}{d\Omega} = \left(\frac{d\sigma}{d\Omega}\right)_{\text{Mott}} \left[ \frac{G_E^2(Q^2) + \tau G_M^2(Q^2)}{1 + \tau} + 2\tau G_M^2(Q^2) \tan^2 \frac{\theta}{2} \right]$$

- $G_E$ : Electric form factor
  - $G_M$ : Magnetic form factor
- Measurement of proton size: 0.8 fm
  - Internal structure of nucleons shown as a mean distribution

# Nucleon structure

- Deep Inelastic Scatterin (DIS) Experiment

$$\frac{d^2\sigma}{dQ^2 d\nu} = \sigma_{\text{Mott}} \left[ W_2(Q^2, \nu) + 2W_1(Q^2, \nu) \tan^2 \frac{\theta}{2} \right]$$





Friedman

Kendall

Taylor

- MIT-SLAC experiment (1969, Friedman, Kendall, Taylor)
  - Scattering cross section does not decrease as  $Q^2$  increases
    - Large angle scattering
    - Point-like components in the proton (parton)
  - Scattering with point-like components rather than scattering by the nucleon as a whole





# Quark-Parton Model (QPM)

- Bjorken scaling rule

$$\frac{d^2\sigma}{dQ^2 dx} = \frac{4\pi\alpha^2}{Q^4} \frac{E'}{E} \frac{1}{x} \left[ F_2(Q^2, x) \cos^2 \frac{\theta}{2} + \frac{Q^2}{2x^2 M^2} 2xF_1(Q^2, x) \sin^2 \frac{\theta}{2} \right]$$

- $F_2$  and  $F_1$  are functions of  $x$  only, independent of  $Q^2$
- Dirac scattering: spin 1/2 target like muon

$$\left( \frac{d\sigma}{dQ^2} \right)_{\text{Dirac}} = \frac{4\pi Z^2 \alpha^2}{Q^4} \left( \frac{E'}{E} \right)^2 \left[ \cos^2 \frac{\theta}{2} + \frac{Q^2}{2M^2} \sin^2 \frac{\theta}{2} \right]$$

- Callan-Gross relation

- Parton spin 1/2 as muon

$$F_2 = 2xF_1$$

$$\frac{d^2\sigma}{dQ^2 dx} = \frac{4\pi\alpha^2}{xQ^4} \{1 + (1-y)^2\} F_2(Q^2, x)$$

- DIS is the superposition of elastic scattering with a point-like component (parton) in the proton


- Parton Distribution Function (PDF)

$$F_2 = x \sum_q e_q^2 q(x)$$

- Internal structure of nucleon shown as parton distribution
- $q(x)$ : parton distribution function of quark  $q$

# From QPM to QCD

- Breaking of the scaling rule
  - When measured precisely, the Callan-Gross relation is broken
    - $F_2$  depends on  $Q^2$
  - Gluon presence
- QCD
  - Asymptotic freedom and confinement






Gross

Politzer

Wilczek


# *Nucleon structure*

- Constituent-quark model
  - Quarks with the effective mass (caused by the gluon)
  - Explains the magnetic moment of the nucleons
  - But, the quark spin cannot explain the nucleon spin (“spin puzzle”)
- Quark-gluon model
  - Current quarks and gluon interaction
  - Initial state of high-energy hadron colliders
- Understanding the differences (or gap) of these models
  - Chiral symmetry (breaking)
  - Confinement



# *Nucleon structure*


- Nucleon: the simplest multi-body system for studying dynamics of confined quarks and gluons
- Simple parton picture
  - 1-dimensional picture: in “longitudinal” direction
  - The nucleon consists of incoherent quarks and gluons
  - Described by the parton distribution functions (PDF)



$x$ : Bjorken's  $x$   
"longitudinal"  
momentum fraction  
(1-dimensional picture)

# Mass

- The Higgs mechanism accounts for only  $\sim 1\%$  of the mass of proton.
- The symmetry breaking emerges the mass.



Symmetry breaking of the Higgs field

Confinement of quarks  
Chiral symmetry breaking

# DIS kinematics

- Deep inelastic scattering (DIS) of lepton

$$Q^2 = -(k - k')^2 = -q^2$$



Measure of resolution power

$$x = \frac{Q^2}{2(p \cdot q)}$$

Measure of momentum fraction by struck quark


$$y = \frac{p \cdot q}{p \cdot k}$$

Measure of inelasticity




# *QCD factorization*

Semi-inclusive DIS process



Proton collision process



- Long distance term
  - unpol. & pol. PDFs – partonic structure of the nucleon
  - fragmentation functions
  - determined from experimental data
  - “universal” property of the nucleon – same in each reaction
  - $Q^2$  dependence calculated by the evolution equation of the perturbative QCD
- Short distance term
  - unpol. & pol. partonic cross section – hard interaction of partons
  - calculated by the perturbative QCD – process dependent
  - the first order (next-to-leading order, NLO) corrections are generally indispensable for a firmer theoretical prediction

$$d\hat{\sigma}_{ab}^c = d\hat{\sigma}_{ab}^{c,(0)} + \frac{\alpha_s}{\pi} d\hat{\sigma}_{ab}^{c,(1)} + \dots$$




# Origin of the nucleon spin 1/2

- EMC experiment at CERN

J. Ashman et al., NPB 328, 1 (1989).

$$\int_0^1 dx g_1^p(x) = \frac{1}{2} \left[ \frac{4}{9} \Delta u + \frac{1}{9} \Delta d + \frac{1}{9} \Delta s \right]$$
$$= 0.123 \pm 0.013(\text{stat}) \pm 0.019(\text{syst})$$



- combining with neutron and hyperon decay data

$$\Delta\Sigma = \Delta u + \Delta d + \Delta s = 12 \pm 9(\text{stat}) \pm 14(\text{syst})\%$$

“proton spin puzzle”  
“proton spin crisis”

- total quark spin constitutes a small fraction of the nucleon spin
- integration in  $x = 0 \sim 1$  makes uncertainty
  - more data to cover wider  $x$  region with more precise data necessary

➔ SLAC/CERN/DESY/JLAB experiments

# Spin

- Spin puzzle
  - Origin of the nucleon spin in the quark-gluon structure

$$\frac{1}{2} = \left[ \frac{1}{2} \Delta\Sigma + L_Q \right] + [\Delta g + L_G]$$


$\Delta\Sigma/2$  = Quark contribution to Proton Spin

$L_Q$  = Quark Orbital Ang. Mom


$\Delta g$  = Gluon contribution to Proton Spin

$L_G$  = Gluon Orbital Ang. Mom

- Quark-spin contribution is only 20%-30% of the nucleon spin
- Gluon polarization measurement with polarized DIS at EIC
  - Small Bjorken- $x$  region with QCD evolution (DGLAP equation)




Integrated gluon polarization down to  $x_{\min}$




# Quark-gluon structure

- Establishing new 3-D picture of the nucleon
  - Nucleon puzzles: spin, radius, mass, pressure...
- Gluon saturation at small- $x$ 
  - Color Glass Condensate (CGC)  $\rightarrow$  Quark Gluon Plasma (QGP)
- and more for standard model & beyond, stability of universe...
  - Neutron EDM, Neutron lifetime, Proton lifetime...
- Importance of precise comparison with Lattice QCD




# Quark-gluon structure

- Establishing new 3-D picture of the nucleon
  - Nucleon puzzles: spin, radius, mass, pressure...
- Ultra-precise electron microscope, revealing the origin of mass and spin in three dimensions
- Gluon saturation at small- $x$ 
  - Color Glass Condensate (CGC)  $\rightarrow$  Quark Gluon Plasma (QGP)
- Discovery of emergent high-density gluon state (gluon condensation)
- Importance of precise comparison with theory, including Lattice QCD




# Precision measurement of PDFs

- Semi-Inclusive DIS (SIDIS)
  - Flavor dependence of quark polarization
  - Transverse-momentum dependence (orbital motion)
- TMD distribution function
  - TMD = Transverse Momentum Dependent
  - Quark, antiquark, gluon
  - 3D distribution including transverse momentum
  - Correlation between spin and orbital motion



Sivers function:  
Correlation between nucleon spin and parton transverse momentum

Sivers function at  $x = 0.1$



December 9, 2022

## Leading Twist TMDs


○ Nucleon Spin      ⊙ Quark Spin

		Quark Polarization		
		Un-Polarized (U)	Longitudinally Polarized (L)	Transversely Polarized (T)
Nucleon Polarization	U	$f_1 = \odot$ Unpolarized		$h_1^\perp = \uparrow - \downarrow$ Boer-Mulders
	L		$g_{1L} = \rightarrow - \leftarrow$ Helicity	$h_{1L}^\perp = \nearrow - \searrow$
	T	$f_{1T}^\perp = \uparrow - \downarrow$ Sivers	$g_{1T}^\perp = \rightarrow - \leftarrow$	$h_1 = \uparrow - \downarrow$ Transversity $h_{1T}^\perp = \nearrow - \searrow$


# TMDs at EIC

Sivers function extracted for valence (left) and sea (right) up quarks from (grey) currently available data and (purple) projection at EIC  $\sqrt{s} = 45 \text{ GeV}$ ,  $10 \text{ fb}^{-1}$


Sivers function of valence quark




Sivers function of sea quark



Transverse-momentum ( $k_T$ ) dependence



Access to the gluon TMDs at EIC  $100 \text{ fb}^{-1}$







# TMD and Aharonov-Bohm effect

- Color Aharonov-Bohm effect
  - Pion production in electron-proton or proton-proton scattering with the polarized proton
  - As the Aharonov-Bohm effect is the left-right asymmetry of the final electron, there is also a left-right asymmetry in the produced pion
  - Siverson effect = Aharonov-Bohm effect in QCD


(magnetic) AB effect



$\Phi_B = 0$



$\Phi_B \neq 0$



interference pattern gets shifted even though the electron never touches upon  $B$


## Speculations on the Siverson effect

Aharonov-Bohm effect	Siverson effect
<p>electron → <math>B</math> → electron</p> <p>(A-fields)</p>	<p>electron → <math>S</math> → pion</p> <p>(A-fields)</p>
<ul style="list-style-type: none"> <li>• L-R asymmetry</li> <li>• magnetic fields only nonzero in solenoid, vector fields provide the effect</li> <li>• topological effect (1, 2, etc)</li> </ul>	<ul style="list-style-type: none"> <li>• L-R asymmetry</li> <li>• gluon-magnetic field only nonzero in hadron, vector fields provide the effect</li> <li>• topological effect (1, 5/4, etc)</li> </ul>
$\exp \left[ (-i) \oint dx \cdot A \right]$	$f_{1T}^{\perp(1)} \sim i \partial_{\xi}^{\alpha} \mathcal{P} \exp \left[ (-i) \oint dx \cdot A \right]$


The first QCD analogue of the Aharonov-Bohm effect?

# Tomography of the nucleon / nucleus


- EIC = color dipole microscope
  - Exclusive process and diffractive process
  - 3D distribution: transverse spatial distribution



## DVCS (Deeply Virtual Compton Scattering)





## Meson Production



- GPD (Generalized Parton Distribution)
    - Spatial imaging of gluons and quarks = tomography
      - HERA: 1<sup>st</sup> generation
      - EIC: 2<sup>nd</sup> generation (high luminosity, heavy ion, polarization)
    - Orbital angular momentum
      - Ji's sum rule
      - Origin of the nucleon spin
- $$J_q^Z = \frac{1}{2} \sum_q \Delta q + \sum_q L_q = \frac{1}{2} \left( \int_{-1}^1 x dx (H^q + E^q) \right)_{t \rightarrow 0}$$

# 3D structure of the nucleon

- How are quarks and gluons confined inside the nucleon?
  - Bag model
    - gluon radius  $>$  charged radius
  - Constituent quark model
    - gluon radius  $\sim$  charged radius
  - Lattice gauge theory (with slow moving quarks)
    - gluon radius  $<$  charged radius
- Need measurement of transverse images of the quarks and gluons in the nucleon
- Proton tomography with **GPD** measurement
  - $R = 0.6 - 0.7$  fm for gluon (HERA) and sea quark (COMPASS)
  - Smaller than 0.85 fm with EM interaction




# Proton radius puzzle

## Conclusions

SpinPhysics@Matsue  
Feb. 23-24, 2021

## Proton Charge Radius Puzzle ??



- disagreements : not yet understood.
- the “correct” proton radius is important.
- further experimental and theoretical efforts.


e-scattering : PRad (JLAB), ULQ2 (Tohoku), MESA (Mainz)  
 $\mu$ -scattering : MUSE (PSI), COMPASS (CERN)

Slide by Suda


# Tomography of the nucleon / nucleus

- DVCS

- Deeply virtual Compton scattering





Spatial distribution of sea quarks at EIC  
 $100 \text{ fb}^{-1}$  and corresponding density of partons in the transverse plane



- Meson production

- Gluon tomography by measuring  $J/\psi$ ,  $\phi$ ,  $\rho$ , etc.
- Precision measurement at large radius with high luminosity


x-dependence of spatial distribution of gluons to be obtained by the exclusive  $J/\psi$  production at EIC





# Mass of the nucleon

- Energy Momentum Tensor Form Factor



## Origin of nucleon mass

Nucleon mass:  $M = \langle p | H | p \rangle$ ,  $H = \int d^3x T^{00}(x)$

Energy-momentum tensor:

$$T^{\mu\nu}(x) = \frac{1}{2} \bar{q}(x) i \vec{D}^{(\mu} \gamma^{\nu)} q(x) + \frac{1}{4} g^{\mu\nu} F^2(x) - F^{\mu\alpha}(x) F^{\nu}_{\alpha}(x)$$

We need theoretical and experimental efforts to decompose nucleon mass for finding its origin.

$$T^{\mu\nu} = \hat{T}^{\mu\nu} + \bar{T}^{\mu\nu} = \left( T^{\mu\nu} - \frac{1}{4} g^{\mu\nu} T^{\alpha}_{\alpha} \right)_{\text{traceless}} + \left( -\frac{1}{4} g^{\mu\nu} T^{\alpha}_{\alpha} \right)_{\text{trace}}, \quad T^{\alpha}_{\alpha} = \bar{q} m q + \frac{\beta(g)}{2g} F^2$$

$H = H_q$  (quark energy) +  $H_g$  (gluon energy) +  $H_m$  (quark mass) +  $H_a$  (trace anomaly)


$$H_q = \int d^3x \bar{q}(x) (-i \vec{D} \cdot \vec{\alpha}) q(x), \quad H_g = \int d^3x \frac{1}{2} (\vec{E}^2 + \vec{B}^2)$$

$$H_m = \int d^3x \bar{q}(x) m q(x), \quad H_s = \int d^3x \frac{9\alpha_s}{16\pi} (\vec{E}^2 + \vec{B}^2)$$


Recent progress on trace-anomaly, gravitational form factor, scale dependence in perturbative QCD:

Y. Hatta, A. Rajan, and K. Tanaka, JHEP 12 (2018) 008;

K. Tanaka, JHEP 01 (2019) 120.



X. Ji, PRL 74 (1995) 1071.




Slide by Kumano




# Mass of the nucleon

- Gravity
  - Calculated from energy and momentum measurements
    - $M^2 = E^2 - p^2$
  - Mass of the nucleon =  $\sim 1$  GeV
- Electroweak
  - Given by the Higgs mechanism
    - $\sim 10$  MeV as the sum of the masses of the 3 valence quarks
  - Cannot be measured in EW
- QCD
  - Measuring the form factor of the Energy-Momentum Tensor (EMT) of QCD
    - Quark and gluon kinetic energy and chiral symmetry breaking (quark and gluon condensations)
  - Can be measured cleanly using spin-2 graviton, but not possible
  - Can be taken out of GPD to be measured by EIC



(c) gravitational



# Mass of the nucleon

- Sum rule for the nucleon mass

Relativistic Motion

Chiral  
Symmetry  
Breaking

Quantum  
Fluctuations

$$M = E_q + E_g + \chi m_q + T_g$$

X. Ji, PRL 74 1071 (1995)


Quark Energy

Gluon Energy

Quark Mass


Trace Anomaly

- How to determine the different contribution not yet reached
- Lattice QCD calculation
  - arXiv:1710.09011, update by K.-F. Liu et al.
- Precision comparison of experiment and theory in the future
  - Mass, spin, pressure, radius,...



# Generalization of the form factor


- Energy Momentum Tensor (EMT)



$$T^{\mu\nu} = \begin{bmatrix} \text{Energy density} & \text{Momentum density} & & \\ T^{00} & T^{01} & T^{02} & T^{03} \\ T^{10} & T^{11} & T^{12} & T^{13} \\ T^{20} & T^{21} & T^{22} & T^{23} \\ T^{30} & T^{31} & T^{32} & T^{33} \\ \text{Energy flux} & \text{Momentum flux} & & \end{bmatrix}$$

Shear stress  
Normal stress (pressure)

- GPD measurement → 3D distribution of mass, spin, pressure, etc. in the proton
  - 1<sup>st</sup> measurement of pressure in the proton using DVCS data from JLab



*Nature*, 557, May 17, 2018

# Nucleon pressure

## Nucleon pressure

$$\langle N(p') | T_q^{\mu\nu}(0) | N(p) \rangle = \bar{u}(p') \left[ A \gamma^{(\mu} \bar{P}^{\nu)} + B \frac{\bar{P}^{(\mu} i \sigma^{\nu)\alpha} \Delta_\alpha}{2M} + D \frac{\Delta^\mu \Delta^\nu - g^{\mu\nu} \Delta^2}{M} + \bar{C} M g^{\mu\nu} \right] u(p)$$

### Recent progress

V. D. Burkert, L. Elouadrhiri, and F. X. Girod,


Nature 557 (2018) 396;

M. V. Polyakov and P. Schweitzer,

Int. J. Mod. Phys. A 33 (2018) 1830025;

C. Lorce, H. Moutarde, and A. P. Tranwinski,

Eur. Phys. J. C 79 (2019) 89.




**Highest pressure in nature** 1 Pa (Pascal) = 1 N/m<sup>2</sup>



Earth atmosphere  
10<sup>5</sup> Pa = 1000 hPa




Center of earth  
10<sup>11</sup> Pa = 100 GPa



Center of Sun  
10<sup>16</sup> Pa = 10 PPa





Neutron star  
10<sup>34</sup> Pa



Hadron  
10<sup>35</sup> Pa

# *Gluon saturation in $e+A$ collisions*




- pQCD and DGLAP & BFKL evolution works with high precision
- Issues with linear DGLAP/BFKL at low- $x$ 
  - Gluon PDF rapid rise violates unitary bound
- New approach: non-linear evolution
  - At very high energy, recombination compensates gluon splitting
  - BK/JIMWLK non-linear effects
  - Saturation characterized by  $Q_s(x)$
  - Describe physics at low- $x$  and low-moderate  $Q^2$






# Gluon saturation in $e+A$ collisions

- Gluon emission
  - Divergence at small  $x$
- Gluon recombination
  - Restriction of divergence
- Gluon saturation in balanced
  - Based on classical idea of the saturation
- Non-linear evolution
  - Saturation of gluon densities characterized by scale  $Q_s(x)$
- Enhancement of  $Q_s$  with  $A$ : saturation regime reached at significantly lower energy in nuclei





$$(Q_s^A)^2 \approx cQ_0^2 \left[ \frac{A}{x} \right]^{1/3}$$



# *Gluon saturation in $e+A$ collisions*


- Inclusive DIS
- Probed by the change of the nuclear structure functions
- Ratio of the structure function  $F_2$ 
  - How quark / gluon distribution and interaction affected in the nucleus?
  - Fermi motion, EMC effect, shadowing, saturation







# *Gluon saturation in $e+A$ collisions*

- Nuclear PDF (nPDF)
  - nPDF measurements for sea quarks and gluons that cannot be reached by LHC and RHIC
  - Discovery of gluon saturation in a small Bjorken- $x$  region



# *Gluon saturation in $e+A$ collisions*



- Semi-inclusive DIS
- Di-hadron correlation
  - Sensitive to the transverse momentum dependence of the gluon distribution and gluon correlations



# *Gluon saturation in e+A collisions*

- Diffractive cross section
  - Most sensitive way to study the gluon saturation
- 10-15% diffractive at HERA e+p
- 25-30% diffractive predicted by CGC at EIC e+A




$$\sigma_{\text{diff}} \propto [g(x, Q^2)]^2$$



# *Gluon saturation in $e+A$ collisions*



- Exclusive vector meson production
  - Momentum transfer  $t$  dependence translated to the transverse spatial distribution of gluons in the nucleus
- Incoherent process (nucleus breaks up)
  - Spatial density fluctuation in nucleus
  - Much larger than the coherent process
- Coherent process (nucleus remains intact)
  - Sensitive to the gluon saturation
  - Identify & veto breakup of the excited nucleus

Meson Production




# *Gluon saturation in e+A collisions*

- What are the emergent properties of dense system of gluons?
  - The gluon saturation describes a new state of matter at extreme high density
- Gluon density saturated where gluon emission and recombination comparable
- First observation of a quantum collective gluonic system




# Gluon saturation

- Precision comparison of experiment and Color Glass Condensate (CGC) as a theoretical model of the gluon saturation
  - Not understandable classically if not discovered?



"Color Glass Condensate"



- Precision understanding of nucleus with the quark-gluon picture necessary as the initial state of the QGP for understanding its production mechanism





# Gluon saturation in $e+A$ collisions



- What are the emergent properties of dense system of gluons?
  - The gluon saturation describes a new state of matter at extreme high density
- First observation of a quantum collective gluonic system
- Precision comparison of experiment and Color Glass Condensate (CGC) as a theoretical model of the gluon saturation
- Precision understanding of nucleus with the quark-gluon picture necessary as the initial state of the QGP for understanding its production mechanism






# Hadronization in the nucleus

- Hadron and jet production from quarks and gluons in the nucleus (cold nuclear matter)
  - Response of nuclear matter to fast moving color charge passing through it?
  - Structure of jet?
- Mass dependence of hadronization
  - Energy loss by light vs. heavy quarks
- Comparison with hot nuclear matter (QGP)



# *EIC physics vs luminosity & energy*



# *Development of lattice QCD*

- Lattice QCD over the next decade will match or exceed experimental accuracy
  - Advances in computational technology
  - Need for computational projects
- Quark and gluon physics advances toward EIC as lattice QCD advances
- Study QCD by comparing precise theoretical calculations with precise experimental measurements to establish an understanding of nucleons, nuclei, and QGP




Supercomputer Fugaku

# *Other physics at EIC*

- Hadron spectroscopy
  - Exotics
- Tensor charge of the nucleon
  - Transversity measurement
- Polarized  $e + d/{}^3\text{He}$  collisions
  - Polarized structure of the neutron
  - “n+p” wave function of the deuteron
- Short range correlations
  - EMC effect by high-momentum “n+p” pairs in nuclei
- High-energy cosmic-ray/neutrino reaction
  - Energy flow in the very forward region
  - Event generator for shower evolution

# *EIC Users Group*

- Formally established in 2016
- More than 1,300 members
  - 36 countries, 266 institutions
  - Experiment (detector, data collection and analysis), theory, computer, accelerator
  - North America 59%, Europe 25%, Asia 12%
- 2020: Yellow report (physics and detector design report) by EIC User Group
- 2020.11: Call for Expressions of Interest (EOI) from the EIC project regarding cooperation in the EIC experimental program
  - EIC-Japan group submitted one EOI from Japan
  - 47 EOIs submitted in total




# *EIC status*

- 2021: Detector collaboration formation and proposal
  - 2021.3: Call for detector proposals
  - 2021.12: Detector proposals due
- 2021.6: CD-1 approval
  - Authorization to begin the project execution phase, starting with preliminary design
  - Cost range \$1.7B - \$2.8B
- 2022.3: Selection of project detector
- 2024: CD-2/3A (performance baseline)
- 2025: CD-3 (start of construction)
- 2032: CD-4A (start of operations)






# EIC status: Schedule

- 2032 CD-4A: Start of operation





# Call for detector proposals

- 3 detector proto-proposals
  - ATHENA
    - A Totally Hermetic Electron-Nucleus Apparatus
    - <https://sites.temple.edu/eicatip6>
  - CORE
    - a Compact detector for the EIC
    - <https://eic.jlab.org/core>
  - ECCE
    - EIC Comprehensive Chromodynamics Experiment
    - <https://www.ecce-eic.org>
- A White Paper for a dedicated IR/detector for lower CM energy being prepared
  - <https://indico.bnl.gov/event/11669>
  - Second PSQ@EIC meeting co-hosted by APCTP



# EPIC detector collaboration


- 2021.3: Call for detector proposal from the EIC project
- 2021.12: Submission of 3 detector proposals
  - EIC-Japan group participates in the ECCE detector consortium
- 2022.3: DPAP (Detector Proposal Advisory Panel) adopts the ECCE detector as the baseline design for the project detector
- EPIC detector collaboration
  - ECCE takes the lead in integrating other detector collaborations
- EIC-Japan group
  - 2020.5: EIC R&D program proposal “Developing a high resolution ZDC for the EIC” (eRD27)
  - ECCE/EPIC ZDC designed by Shimizu




ECCE/EPIC ZDC

# *EIC Interaction Region (IP6)*

- Extensive integration of forward and backward detector elements into the accelerator lattice



# EIC Far-forward region



Detector	(x,z) Position [m]	Dimensions	$\theta$ [mrad]	Notes
ZDC	(-0.96, 37.5)	(60cm, 60cm, 1.62m)	$\theta < 5.5$	$\sim 4.0$ mrad at $\phi = \pi$
Roman Pots (2 stations)	(-0.83, 26.0) (-0.92, 28.0)	(30cm, 10cm)	$0.0 < \theta < 5.5$	$10\sigma$ cut.
Off-Momentum Detector	(-1.62, 34.5), (-1.71, 36.5)	(50cm, 35cm)	$0.0 < \theta < 5.0$	$0.4 < x_L < 0.6$
B0 Trackers and Calorimeter	(x = -0.15, $5.8 < z < 7.0$ )	(32cm, 38m)	$6.0 < \theta < 22.5$	$\sim 20$ mrad at $\phi=0$





# *EIC Japan*

- 2019: Master Plan 2020 proposal of Electron-Ion Collider (EIC)
  - Selected as a major academic research project
  - Core institutions: Yamagata Univ. & RIKEN
  - Participating institutions: Kobe Univ., Nihon Univ., KEK, etc.
- Collaboration including nuclear-physics community and high-energy community
  - Nuclear physics: Yamagata U., RIKEN, Nihon U., U. of Tsukuba, JAEA
  - High-energy physics: Kobe U., Shinshu U., Kyushu U., KEK
  - Cosmic ray: Nagoya U., ICRR
- 2020.5: EIC detector R&D program eRD27
  - “Developing a high resolution ZDC for the EIC”
  - Collaboration with Kansas U., ODU, etc.
- 2020.11: Expression of interest (EOI) from EIC-Japan
  - Forward hadron calorimeter
    - Cooperation with UCLA & Korean group
  - Zero-degree calorimeter (EM & hadron)
    - Cooperation with eRD27
  - Silicon detector
    - Cooperation with ANL, BNL, etc.



# EIC Japan

- 2022: Science Council of Japan, "Medium- and Long-term Research Strategy for Science"
  - 7/18 Status report at future planning committee meeting of Nuclear Physics Committee
  - Participation in the EIC project as part of the High Energy QCD Frontier Initiative
- Mailing list
  - [eic-japan-l@ml.riken.jp](mailto:eic-japan-l@ml.riken.jp)
- Meeting held on Thursdays at 10:30AM
  - Participating groups: RIKEN, Yamagata Univ., JAEA, Nihon Univ., Virginia Univ., Kobe Univ., Shinshu Univ., Kyushu Univ., KEK, Nagoya Univ., Univ. of Tokyo ICRR, Univ. of Tsukuba, Tsukuba Univ. of Technology, Univ. of Tokyo CNS, Hiroshima Univ., Nara Women's Univ.
- EIC detector prototype R&D
  - ZDC
  - AC-LGAD




# *EIC-Japan group involvement*


- 2015.4: EIC Letter of Interest from Asian countries
  - 20 participants from Japan: RIKEN, Yamagata, Tokyo Tech, Juntendo, KEK, Kyorin, Kyoto, Niigata, Tohoku, Tokyo Science
  - 7 from China, 3 from India, 4 from Korea
  - To support EIC for NSAC Long Range Plan 2015
- 2019: Science Council of Japan Master Plan 2020 proposal of EIC
  - Collaboration including nuclear-physics community and high-energy community
  - Core institutions: Yamagata and RIKEN
  - Participating institutions: Kobe, Nihon, KEK, etc.
- 2020: Yellow Report
- 2020.5: eRD27 “developing a high resolution ZDC for the EIC”
- 2020.11: Expression of Interest (EOI) from EIC-Japan
- 2021.3-12: Call for detector proposal from the EIC project
  - EIC-Japan group participates in the ECCE detector consortium
- 2022: Science Council of Japan “Medium- and Long-term Research Strategy for Science”
  - EIC project proposal to be submitted as a part of the High-Energy QCD Frontier Initiative

# *Interest in contributing to ZDC*



- ECCE/EPIC ZDC design
  - Simulation
  - Performance evaluation
- ALICE-FoCal-E technology: Tungsten/Silicon
  - Test beam studies ongoing
- Radiation tolerance test by neutron irradiation
- RIKEN, Tsukuba, Tsukuba Tech, Kobe, Shinshu, Yamagata, JAEA, Nihon, Kyushu, KEK, Nagoya, Tokyo ICRR



ECCE/EPIC ZDC




ALICE FoCal-E  
R&D






Neutron irradiation at  
RIKEN RANS

# *Interest in contributing to (AC-)LGAD Barrel*

- Construction of (AC-)LGAD Barrel based on our past experience of PHENIX VTX silicon detector construction and present experience of sPHENIX INTT silicon detector construction
- HPK LGAD development by KEK group
  - To be combined with some readout ASIC
- RIKEN, Hiroshima, Nara Women's, Tokyo CNS, Kyushu, KEK




sPHENIX INTT construction



HPK LGAD development

# Calorimeter development

- ALICE-FoCal
  - 2027-29 LHC Run-4
  - FoCal-E Pad & Pixel
    - Pad detector led by Univ. of Tsukuba group (and Indian group)
    - Pixel detector led by European group
  - Test beam activities ongoing
    - Pad detector at ELPH, Tohoku Univ.
    - Total system at CERN SPS
  - TRD to be made in 2022





	19	2020				2021				2022				2023				2024				2025				2026				2027			
	Q4	Q1	Q2	Q3	Q4	Q1	Q2	Q3	Q4	Q1	Q2	Q3	Q4	Q1	Q2	Q3	Q4	Q1	Q2	Q3	Q4	Q1	Q2	Q3	Q4	Q1	Q2	Q3	Q4	Q1	Q2	Q3	Q4
LHC		LS2				Run-3												LS3								Run-4							
LoI		R&D																															
R&D		R&D				R&D				R&D				R&D				R&D				R&D				R&D							
Test beam						Test beam				Test beam				Test beam				Test beam				Test beam				Test beam							
TDR						TDR				TDR				TDR				TDR				TDR				TDR							
Final design						Final design				Final design				Final design				Final design				Final design				Final design							
Production, construction, test of module										Production, construction, test of module				Production, construction, test of module				Production, construction, test of module				Production, construction, test of module				Production, construction, test of module							
Pre-assembly, calibration with test beam														Pre-assembly, calibration with test beam				Pre-assembly, calibration with test beam				Pre-assembly, calibration with test beam				Pre-assembly, calibration with test beam							
Installation and commissioning																						Installation and commissioning				Installation and commissioning							
Physics data taking																										Physics data taking							



# Irradiation test

- Measurement of the radiation hardness of the ALICE-FoCal-E Pad sensors
  - To determine if the sensor is sufficiently radiation hard to radiation dose/fluence at zero degree of EIC
  - At ALICE-FoCal, 1-MeV neutron equivalent fluence  $\Phi_{eq} < 10^{13} \text{ n}_{eq}/\text{cm}^2$ , or Total ionization dose (TID) of 1.5 kGy
- HPK Pad sensor
  - Test its baby chip and MPD
- Options: p-type or n-type
  - Type inversion from n-type to p-type at  $10^{12} \text{ neutron}/\text{cm}^2$







# EIC-ZDC

- More than  $2 \times 10^{13}$  neutron/cm<sup>2</sup> in one year at EIC-ZDC
  - More than 10 times higher radiation than ALICE-FoCal

Electron-Proton collisions. IP6, p(275)+e(10). Si lifetime in ZDC.



Assume the ep - collision rate is  $1.E+6$  [ep/s]

$$\text{Critical rate } \frac{dN_{ep+pg}}{dt} = 8.E - 1 \left[ \frac{N}{cm^2/ep} \right] * 1.E+6 [ep/s] + \frac{dN_{pg}}{dt} = 8.E+5 [Hz/cm^2] + .4 E+5 [Hz/cm^2]$$

$$\text{ZDC Silicon LifeTime} = \frac{1.E+14 [1/cm^2]}{8.4E+5 [Hz/cm^2]} = \sim 0.12 E+9 [s] = \sim 4 \text{ years.}$$


# Test plan & menu

- Proton 7MeV, 100 $\mu$ A, 6 x 10<sup>13</sup> proton/s
  - Maximum current stable produced about 40 $\mu$ A
- Neutron 5MeV max, 10<sup>12</sup> neutron/s from the target
- 2 cm from the target: 10<sup>8</sup> neutron/cm<sup>2</sup>/s
  - After the Be target, several x 10 $\mu$ m silver wax, 5mm vanadium, 5mm water, and 5mm titanium
- Maximum operation time per day about 5 hours
  - Up to 1 hour of continuous operation with a 10-minute stop for the next irradiation
- Measurement of I-V and C-V characteristics every hour to evaluate  $U_{\text{dep}}$



位置	1時間	2時間	3時間	4時間	5時間
2cm	0.36	0.72	1.08	1.44	1.80
5cm	0.06	0.12	0.17	0.23	0.29
10cm	0.01	0.03	0.04	0.06	0.07


Table 1: 1日の運転で得られる照射量の見積もり ( $\times 10^{12} n_{\text{eq}}/\text{cm}^2$ ).



# EIC Detector R&D

- “Developing a High Resolution ZDC for EIC”

- Submitted in 2020.5
- eRD27 approved in 2020.8



- Soft photon detection

- Large aperture
- Full absorption calorimeter

- EM and hadron calorimeter

- Acceptance
- Energy, position and  $p_T$  resolution
- ALICE-FoCal R&D by RIKEN & Tsukuba


- Radiation hardness

## Performance requirements and resources requested


Detector R&D	Physics	Performance requirements	Resource requested	Support & collaboration
Soft photon detection	e+A nuclear breakup veto	$E_\gamma \leq 300$ MeV	detector simulation	This proposal Calorimeter consortium
		acceptance	acceptance simulation	This proposal BeAGLE group
		detector technology	detector R&D	N/A in FY21
EM + hadron calorimeter	e+A collision geometry	neutron multiplicity	high resolution not necessary	BeAGLE group
	spectator tagging	energy & position resolution	detector simulation	This proposal
	meson structure	neutron & $\Lambda$ acceptance	detector simulation	This proposal Meson structure WG
	detector technology		FoCal R&D LHC-ZDC R&D	RIKEN Kansas Univ.
	calibration scheme		design & simulation system test	This proposal N/A in FY21
Radiation hardness		radiation dose	simulation study	This proposal Kobe Univ.
		detector technology	radiation test	This proposal Calorimeter consortium

# Physics at zero degree of EIC

- Spectator tagging
  - Neutron structure
    - Neutron spin structure, S & D waves
  - Nucleon interactions
    - SRC/EMC at large  $x$
    - Diffraction and shadowing at small  $x$
- Geometry tagging in  $e+A$  collisions
  - $b$ : impact parameter &  $d$ : path length
  - “Centrality” (high  $d$ ) & “Skin” (low  $d$ )
  - Breakup determination & veto with ZD
    - + forward photons requiring wide aperture
- Event generator




- GCF: [https://www.mit.edu/~src\\_emc/fri/schmidt\\_20190322.pdf](https://www.mit.edu/~src_emc/fri/schmidt_20190322.pdf)
- BeAGLE: <https://wiki.bnl.gov/eic/index.php/BeAGLE>



# *$e + A$ collision at zero degree*

- Breakup of the excited nucleus
  - Evaporated neutrons (& protons)
    - Separate the coherent process  $\sim 90\%$
  - Photons from de-excitation of the excited nucleus
  - Requirement to measure neutrons and photons at zero degree in a wide  $t$  range
- Event-by-event characterization of collision geometry
  - Tagged through forward neutron multiplicities at zero degree
  - $b$ : impact parameter
  - $d$ : path length of struck parton in nucleus
  - “centrality” (high  $d$ ) & “skin” (low  $d$ )
  - Study of nuclear medium effects






# *e + A collision at zero degree*

Slide by C. Hyde

## ZDC EMCAL: DEEP EXCLUSIVE NUCLEI

- Gluon Density from e.g.  $^{208}\text{Pb}(e,e'\phi)^{208}\text{Pb}$ 
  - Final state nucleus is lost in beam envelope
  - Veto breakup of Pb nucleus.
  - Thousands of bound states excitable by photo-excitation
    - These will wash out diffractive minima.
    - Possible veto by detection of boosted decay photons
    - At  $P_{\text{Pb}} = 275 \cdot Z \text{ GeV}$ , boost  $\gamma = 117$
    - Each photon has 32% detection probability within 4mr cone




- Removing excited nucleus event by detecting excitation photon
- Soft photon  $\sim 300 \text{ MeV}$
- Low detection probability within 4 mrad cone



# $e + d/{}^3\text{He}$ collision at zero degree


- Spectator tagging
  - Neutron structure
    - Neutron spin structure, S & D waves
  - Nucleon interactions
    - Short-range correlation (SRC) and EMC effect at large  $x$
    - Diffraction and shadowing at small  $x$



# Physics at zero degree of EIC

- Short range correlation (SRC)
  - ~20% of nucleons in SRC pairs
    - 18% p-n pairs
    - Large relative momentum ( $> 300$  MeV/ $c$ )
    - Small c.m. momentum and spatially very close each other
  - EMC effect
    - Nuclear structure modification found in nuclear DIS in the EMC experiment
    - Nuclear PDF significantly modified by SRC pairs
- Tagged DIS at JLab  $\rightarrow$  EIC
  - e+D at JLab: Hall B & C
  - e+D & e+A at EIC
- Tagged SRC at EIC

## Nucleon Momentum Distribution




OLD DOMINION

Hauenstein | 09/24/2019

8

## EMC and SRC Correlation



Weinstein et al., PRL 106, 052301 (2011), Hen et al., PRC 85, 047301(2012)

Hauenstein | 09/24/2019


14

slides by Heuenstein

# Spectator tagging

## ZDC RESOLUTION: SINGLE NEUTRON EVENTS

- Measuring the properties of a bound proton:  
Spectator tagging: e.g.  $D(e,e'n)X$ 
  - $P_D = 275 \text{ GeV}/c \rightarrow p_n = P_D(1+\alpha)/2 \approx 137 \text{ GeV}/c$ 
    - Rest frame neutron momentum  $\approx \alpha M$
  - If ZDC resolution = 50%  $[\text{GeV}/E_n]^{1/2}$   
 $\rightarrow 4.5\% \text{ @ } 137 \text{ GeV}/c$ 
    - $\sigma(\alpha) \approx \sigma(p)/p \approx 0.045$   
 $\rightarrow \text{Rest-frame } \sigma(p_n) \approx 40 \text{ MeV}/c$
- Spatial resolution 1 cm ?
  - $\sigma(p_T) \approx (137 \text{ GeV}/c) (1 \text{ cm})/(32m) = 43 \text{ MeV}/c$





Slide by C. Hyde

- $p_T$  resolution equivalent to beam spread  $\sim 40\text{-}50 \text{ MeV}/c$
- Spatial resolution 1cm  $\rightarrow p_T$  resolution  $\sim 40 \text{ MeV}/c$
- ZDC energy resolution  $50\%/\sqrt{E} \text{ (GeV)}$  or  $4.5\% \text{ @ } 137 \text{ GeV}/c$   
 $\rightarrow p_T$  resolution  $\sim 40 \text{ MeV}/c$

# Far-forward physics at EIC

- Mass of the proton, pion, kaon
  - Visible world mainly made of light quarks: its mass emerges from quark-gluon interactions, Higgs mechanism hardly plays a role
  - Strange quark is at the boundary: both emergent-mass and Higgs-mass generation mechanism are important
- Proton
  - EIC will allow determination of an important term contributing to the proton mass, the so-called “QCD trace anomaly”
- Pion and kaon
  - EIC will allow determination of the quark and gluon contribution to mass with the Sullivan process


## Sullivan process




# Physics at zero degree of EIC

- Leading baryons
  - Fragmentation
  - One pion exchange (OPE)

One Pion Exchange (OPE)



## LN in DIS






$$d\sigma_{\gamma^* p \rightarrow nX} = f_{\pi/p}(x_L, t) \times d\sigma_{\gamma^* \pi \rightarrow X}$$

The distribution of  $p_T^2 (=t)$  is defined solely by the pion flux

Sensitivity to the pion flux

$p_T^2$  dependence in bins of  $x_L$



Slope of exponential  $p_T^2$  dependence computed to various pion-flux models

18


slide by Ciesielski

Inconsistency @ HERA

→ Need more data to understand production mechanism

# EIC IR design

- Acceptance
  - 25 mrad crossing angle for EIC at BNL
  - Forward magnet aperture  $\pm 4$  mrad opening angle for ZDC
- Sufficient transverse size to avoid transverse leakage
  - $\sim 2$  interaction length
  - e.g. 60cm x 60cm



IR design of EIC at BNL



# *EICUG Yellow Report*

- ~400 authors / ~150 institutions / ~900 pages with strong international contributions
  - Volume I: Executive Summary
  - Volume II: Physics
  - Volume III: Detector
- <https://arxiv.org/abs/2103.05419>






**EIC YELLOW REPORT**  
Volume I: Executive Summary



**EIC YELLOW REPORT**  
Volume II: Physics





**EIC YELLOW REPORT**  
Volume III: Detector



# Call for detector proposals

- Detector 1 (D1)
  - Within the scope of the EIC project
- Detector 2 (D2)
  - Not within the scope of the EIC project
  - How to realize it being explored
- Location of D1 and D2 between IP6 and IP8 is left open, so far assumed D1 will go to IP6




# *Asian collaboration for EIC*

- 2015.4 EIC Letter of Interest from Asian countries
  - 20 from Japan
    - RIKEN, Yamagata U, Tokyo Tech, Juntendo U, KEK, Kyorin U, Kyoto U, Niigata U, Tohoku U, Tokyo U of Science
  - 7 from China
    - CIAE, IMP, Nanjing U
  - 3 from India
    - TIFR, NISER
  - 4 from Korea
    - Seoul National U, Korea U, Daegu U, Chonbuk National U
  - To support EIC for NSAC Long Range Plan 2015
- 2020.11 Eols from Asian countries
  - EIC-Japan Expression of Interest
  - Eol for China
  - Eol for Indian Consortium
  - Eol for Korean Institutions

## Letter of Interest Participation in the US Electron-Ion Collider (EIC) from Asian countries


With this letter we want to express our interest in participating in the US EIC project. The EIC project being discussed in the Long Range Plan process of the NSAC is the most promising project in the world to be realized in a timely manner. It is a new collider which will be able to collide polarized electrons with polarized protons or nuclei. We will be able to have 100-1,000 times higher luminosity per nucleon than HERA. It promises to lead to deep understanding of high-energy QCD and the development of a novel physics field based on QCD where the gluon plays a leading role. The mass of the nucleon and the nuclei originates from gluon interactions and dynamics, and the confinement of the quarks inside the nucleon is caused by the gluons. We are keenly interested in this science, and want to strongly support the US EIC project, through a long-term collaboration for investigations of the novel gluon related physics at EIC.

# ECCE tracking system



## • MAPS

- 3-layer silicon vertex, 2-layer silicon sagitta, 5 disks in the hadron endcap, 4 disks in the electron endcap
- For primary and secondary vertex reconstruction
- Low material budget: 0.05%  $X/X_0$  per layer
- High spatial resolution: 10  $\mu\text{m}$  pitch MAPS (ALICE ITS3)
- TowerJazz 65 nm technology (ongoing R&D Si Consortium)




## • AC-LGAD


- TOF layer integrated with PID




# ECCE charged particle ID



- Need to separate:
  - Electrons from photons
  - Electrons from charged hadrons
    - Calorimeter
  - Charged pions, kaons and protons from each other
    - TOF and Cherenkov
- AC-LGAD based TOF system
  - Hadron PID in momentum range below the thresholds of the Cherenkov detectors



# ECCE far-forward region





Detector	(x,z) Position [m]	Dimensions	$\theta$ [mrad]	Notes
ZDC	(-0.96, 37.5)	(60cm, 60cm, 1.62m)	$\theta < 5.5$	$\sim 4.0$ mrad at $\phi = \pi$
Roman Pots (2 stations)	(-0.83, 26.0) (-0.92, 28.0)	(30cm, 10cm)	$0.0 < \theta < 5.5$	$10\sigma$ cut.
Off-Momentum Detector	(-1.62, 34.5), (-1.71, 36.5)	(50cm, 35cm)	$0.0 < \theta < 5.0$	$0.4 < x_L < 0.6$
B0 Trackers and Calorimeter	(x = -0.15, $5.8 < z < 7.0$ )	(32cm, 38m)	$6.0 < \theta < 22.5$	$\sim 20$ mrad at $\phi=0$




# *ECCE far-forward region*

- B0 spectrometer
  - 4 AC-LGAD tracers with 30 cm spacing between each layer providing charged particle detection for  $6 < \theta < 22.5$  mrad
- Roman pots
  - Two double-layer  $25 \times 12$  cm<sup>2</sup> AC-LGAD stations
  - Located inside the beam line
  - $10 \sigma$  from the main beam
- Off-momentum detectors
  - AC-LGAD for fast timing
- Zero-Degree Calorimeter (ZDC)
  - W/Si EMCal
    - Position &  $p_T$  resolution
    - Radiation hardness
  - Pb/Si HCal



# *ECCE far-backward region*

- Electron-going region
- Low  $Q^2$  tagger
  - Double-layer AC-LGAD tracker
  - 40.5 cm x 40.5 cm at 24 m and 30 cm x 30 cm at 37 m from IP
- Luminosity monitor
  - AC-LGAD and PWO
  - Accuracy of the order of 1% or relative luminosity determination exceeding  $10^{-4}$  precision



# *Summary of this talk*

- Physics at EIC
  - Origin of nucleon mass and spin
  - 3D structure of the nucleon and nucleus
  - Gluon saturation
  - Hadronization
  - Ultra-precise electron microscope, revealing the origin of mass and spin in three dimensions.
  - Discovery of emergent high-density gluon state (gluon condensation)
- Physics at zero-degree
  - e+A breakup of the excited nucleus & collision geometry
  - e+d/<sup>3</sup>He spectator tagging & short-range correlation
  - pion and kaon structure (and mass)
- EIC status
  - EIC Users Group
  - EIC-Japan
  - EPIC detector collaboration
  - The EIC project and EPIC experiment are progressing very well
  - EIC-Japan Group is developing steadily
  - Japan takes the lead in completing QCD research at EIC

# *Summary of this talk*

- Physics at EIC
  - Origin of nucleon mass and spin
  - 3D structure of the nucleon and nucleus
  - Gluon saturation (CGC)
  - Hadronization
  - Ultra-precise electron microscope, revealing the origin of mass and spin in three dimensions.
  - Discovery of emergent high-density gluon state (gluon condensation)
- EIC status
  - EIC Users Group
  - EIC-Japan
  - EPIC detector collaboration
  - The EIC project and EPIC experiment are progressing very well
  - EIC-Japan Group is developing steadily
  - Japan takes the lead in completing QCD research at EIC